

Interaktionen von RNAs und Proteinen



Sonja Prohaska

Computational EvoDevo
Universitaet Leipzig

SS17


(Multi) Protein Complexes

- ▶ two or more associated polypeptide chains
- ▶ **with homologous structure/function**
formation of homo- and heterodimers to oligomers from monomers
e.g. helicase
- ▶ **with different structure/function**
built from (core) subunits and accessory proteins
e.g. PRC2 complex
- ▶ protein complexes are a form of quaternary structure
- ▶ strong bonding is based on disulphid bridges, salt bridges, hydrophobic contacts, electron sharing



Protein-Protein Interactions


- ▶ define interaction interfaces/surfaces
- ▶ bonding is based on hydrophobic contacts, Van der Waals forces, hydrogen bonds



- a** two globular proteins with preformed surfaces
- b** two globular proteins with an induced binding surface
- c** rigid globular protein with a peptide
- d** flexible globular protein with a peptide
- e** interaction of two peptides


Protein-Protein Interaction Domain: SH2 Domain

- ▶ **Src Homology 2**
- ▶ around 100 aa
- ▶ 2 α -helices and 7 β -strands
- ▶ (one large β -sheet)
- ▶ known to identify a sequence of 3-6 aa
- ▶ high affinity to phosphorylated tyrosine
- ▶ function – signaling
- ▶ found in about 100 human proteins




Protein-Protein Interaction Domain: SH3 Domain

- ▶ **Src Homology 3**
- ▶ beta-barrel fold: six β -strands forming two tightly packed anti-parallel β -sheets
- ▶ contacts proline-rich peptide sequence: -X-P-p-X-P- (X – aliphatic amino acid; p – sometimes proline)
- ▶ function – signaling
- ▶ found in about 300 human proteins




Protein-Protein Interaction Domain: SAM Domain

- ▶ **Sterile Alpha Motif**
- ▶ around 70 aa
- ▶ small five-helix bundle
- ▶ seems to possess the ability to bind RNA
- ▶ has two large interfaces
- ▶ can form dimers
- ▶ found in small group of genes




Protein-Protein Interaction Domain: PDZ domain

- ▶ 80-90 aa
- ▶ 5 β -sheets, some helices
- ▶ binds to C-terminus of binding partner
by adding a β -strand to the β -sheet
- ▶ multiple PDZs per protein
increase specificity
- ▶ 260 PDZ in 180 human genes



Yeast Two-Hybrid (Y2H)

- ▶ test if X binds Y
 - express fusion protein X-DB (DB ... DNA binding domain)
 - express fusion protein Y-AD (AD ... activation domain)
 - long linkers between X and DB, and Y and AD



Yeast *Saccharomyces cerevisiae*

- ▶ genome size: 12.5×10^6 bp
- ▶ about 5770 genes
- ▶ about 6100 proteins (about 2000 uncharacterized)
- ▶ 5100 soluble: 47% cytoplasm, 27% nucleus
- ▶ haploid and diploid living forms

High-Throughput Yeast Two-Hybrid screening (HT-Y2H)

- ▶ a *bait* library (e.g. 192 proteins)
- ▶ theoretically any number of bait
- ▶ a *prey* library (e.g. about 6000 proteins)
- ▶ *bait* encodes fusion protein: DB and protein X
- ▶ *prey* encodes fusion protein: AD and protein Y
- ▶ array of *prey*-expressing haploid cells
- ▶ library of *bait*-expressing haploid cells
- ▶ mate *bait* with *prey*
- ▶ retrieve diploid cells expressing reporter gene due to protein-protein interaction (PPI)

It is difficult to sample **all** possible binary combinations of proteins using the library screening methods.

High-Throughput Yeast Two-Hybrid screening (HT-Y2H)

How it works: see blackboard

- ▶ 1-30 positives per *bait*
- ▶ only 20% true positives
- ▶ reason: self activation of reporter gene
- ▶ solution: two independent screens per bait
- ▶ even better solution: varify with other method
- ▶ found 87 baits in 281 PPI
- ▶ very low resolution
- ▶ Problem: construction of artificial fusion proteins



co-Immunoprecipitation (co-IP)

Co-Immunoprecipitation is not a high-throughput method.


It is used for validating PPI predictions, e.g. from HT-Y2H.

- ▶ antibodies against epitope on known protein (e.g. *bait*)
- ▶ pull the entire protein complex (“pull-down”)
- ▶ works if proteins bind to each other tightly
- ▶ Problem: protein might hide epitope in complex
- ▶ Solution: antibody against different epitope on same protein
- ▶ Solution: antibody against proposed binding partner (double-check)
- ▶ Problem: no detection of transient interactions


co-Immunoprecipitation (co-IP)



Real-Time Single-Molecule co-IP




Co-IP
 +
 real-time
 detection
 (Time resolution
 ~50 ms)



Advantage: suited for transient interactions, kinetics studies (time resolution of 50ms)

Disadvantage: requires EGFP labeling of prey

Tandem Affinity Purification (TAP)



Tandem Affinity Purification (TAP)

- ▶ target protein is tagged on C-terminus
- ▶ tag: calmodulin binding peptide (CBP), cleavage site, protein A
- ▶ 1. purification: beads with IgG bind protein A, wash
- ▶ cleave at cleavage site, elute
- ▶ 2. purification: beads with calmodulin bind CBP, wash
- ▶ analyse protein complexes with mass spectrometry

TAP and mass spectroscopy


- ▶ using LC-MS, and MALDI/TOF-MS
- ▶ 4562 different tagged proteins
- ▶ 2708 proteins in 7123 PPI (2006)
- ▶ **improvement!**

Mass Spectrometry - General Idea

Identify and quantify multiple proteins in one run


- ▶ starting with the pool of proteins
- ▶ coarse-grained protein separation by size (electrophoresis)
- ▶ protein fragmentation e.g. with trypsin
- ▶ fine-grained peptide separation with liquid chromatography (LC)
- ▶ electrospray ionization of peptides
- ▶ mass analysis → MS spectrum
- ▶ (collide peptides with neutral gas, even smaller fragments, mass analysis → MS/MS spectrum)
- ▶ **bioinformatic analysis** of MS spectrum

Protein Mass Spectrometry






Mass Spectrometry

- ▶ mass (m) charge (q) ratio
- ▶ **peptide mass fingerprints**
are stored in databases
- ▶ for comparison



- ▶ some amino acids have identical masses
- ▶ utilize a sequence homology search in parallel
- ▶ problem: modifications change mass \rightarrow misidentification

Interactome



Propose “highly significant clustering between essential proteins”

Interactome as a Graph?

- ▶ only binary protein interactions are considered
- ▶ interesting interactions might be transient
- ▶ only a static picture lacking dynamics and context
- ▶ A interacts with B and B interacts with C
 - ▶ at the same time?
 - ▶ in the same compartment or cell type?
 - ▶ Is there direct or indirect interaction of A and C?
 - ▶ Do A, B and C form a complex?
- ▶ would we see pathways?
- ▶ what can we infer from such a network?

Literature

Hong-Won Lee, Ji Young Ryu, Janghyun Yoo, Byungsan Choi, Kipom Kim, Tae-Young Yoon. *Real-time single-molecule coimmunoprecipitation of weak protein-protein interactions*. Nature Protocols 2013; 8, 2045-2060

For further reading:

<https://www.intechopen.com/books/protein-protein-interactions-computational-and-experimental-tools>